

Visit
FREETESTPAPER.com
for more papers



Website: freetestpaper.com



[Facebook.com/freetestpaper](https://www.facebook.com/freetestpaper)



[Twitter.com/freetestpaper](https://www.twitter.com/freetestpaper)

Name

4047/01

16/S4PR2/AM/1

ADDITIONAL MATHEMATICS

PAPER 1

Wednesday

3 August 2016

2 hours



VICTORIA SCHOOL

**PRELIMINARY EXAMINATION TWO
SECONDARY FOUR**

Additional Material: Answer Paper

READ THESE INSTRUCTIONS FIRST

Write your name, class and register number on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use paper clips, highlighters, glue or correction fluid.

Answer **all** questions.

If working is needed for any question it must be shown with the answer.

Omission of essential working will result in loss of marks.

You are expected to use a scientific calculator to evaluate explicit numerical expressions.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place.

For π , use either your calculator value or 3.142, unless the question requires the answer in terms of π .

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 80.

This paper consists of 6 printed pages, including the cover page.

[Turn over

1 (i) Simplify $\left| 21 - 14x \right| - \left| \frac{2}{3}x - 1 \right|$. [2]

(ii) Hence, solve $|21 - 14x| = \left| \frac{2}{3}x - 1 \right| + 40 - 15x$. [3]

2 The range of solutions for x such that $a+bx-4x^2 > 0$ is $-\frac{1}{2} < x < 3$. Find the value of a and of b , where a and b are real numbers. [4]

3 (a) Solve $4^x - 20(4^{-x}) = 1$. [3]

(b) Given that $\frac{5^{4-\frac{x}{2}}}{625(5^x)} = \frac{25^y}{\sqrt{125^x}}$, find the value of $\frac{x}{y}$. [3]


4 In the expansion of $\left(ax + \frac{1}{x}\right)^n$ in descending powers of x , where a and n are positive integers, the fourth term of the expansion is the constant term.

(i) Find the value of n and hence, express the constant term in terms of a . [4]

(ii) Using your value of n in (i), determine if a term in $\frac{1}{x^3}$ in the expansion $(1-x)\left(ax + \frac{1}{x}\right)^n$ exists. [2]

- 5 (a) Solve $\sqrt{1-2x}-3x=1$. [3]

(b)



In the diagram above, ACD is a triangle such that B lies on AC , $AB = (5 - \sqrt{12})$ cm, $CD = (3 + \sqrt{3})$ cm, $\angle BDC = 30^\circ$ and $\angle BCD$ is a right angle. Find AD^2 in the form $p + q\sqrt{3}$, where p and q are constants. [4]

- 6 (a) Differentiate $\ln \sqrt{\frac{1-3x}{e^{-x}}}$. [3]

- (b) Given that $\int_1^4 f(x) dx = 6$, $\int_3^6 f(x) dx = 2$ and $\int_7^4 f(x) dx = -3$, find

(i) $\int_1^7 f(x) dx$, [1]

(ii) $\int_1^3 f(x) dx + \int_6^7 f(x) dx$, [1]

(iii) the value of h , where $\int_1^4 hx^2 + 2f(x) dx = 180$. [2]

- 7 The equation of a curve is $y = 3\left(\frac{x}{4} + a\right)^{\frac{5}{6}}$. The normal to the curve at $x = \frac{1}{2}$ is parallel to the line $5y + 4x = 2$.

(i) Show that $a = -\frac{7}{64}$. [4]

(ii) Find the equation of the tangent to the curve at $x = \frac{1}{2}$. [2]

(iii) Show that the curve is an increasing function for $x > \frac{7}{16}$. [2]

- 8 It is given that $\cos 140^\circ = -k$ and $\tan A = -\frac{3}{4}$, where k is a positive number and A is a reflex angle. Without finding the value of A or of k ,

(i) find the exact value of $\cos 2A$, [2]

(ii) express $\tan 50^\circ$ in terms of k , [3]

(iii) express $\sin(40^\circ + A)$ in terms of k . [3]

- 9 The points $P(1, -2)$ and $Q(1, 4)$ lie on the circumference of a circle with centre C . If the circle is reflected in a vertical line, P and Q remain unchanged in the reflection and the x -coordinate of the centre of the reflected circle is 5.

(i) State the equation of the vertical line of reflection. [1]

(ii) Show that the equation of the circle with centre C is $x^2 + y^2 + 6x - 2y - 15 = 0$. [3]

(iii) The line $3y + 4x = -9$ intersects the circle with centre C at two points, A and B . Find the coordinates of A and of B . [4]

(iv) Determine if AB is a diameter of the circle with centre C . [1]

- 10 A particle moves in a straight line such that at t seconds after passing point O , its velocity v m/s is given by $v = t - 7 + \frac{12}{t+1}$, where $t > 0$.

Find


(i) the acceleration of the particle when it is first instantaneously at rest. [3]

(ii) an expression for the displacement of the particle from O . [3]

(iii) the total distance travelled by the particle from $t = 0$ to $t = 5$. [3]

11 Solutions to this question by accurate drawing will not be accepted.

The diagram shows the quadrilateral $ABCD$ in which point A is $(-2, 1)$ and point C is $(25, 7)$. The point $P\left(7\frac{3}{5}, 7\right)$ lies on AD such that $AP : PD = 3 : 2$. The midpoint of BC , point M , lies on the x -axis and directly below point D .



- (i) Find the coordinates of points D , M and B . [6]
- (ii) Determine if $\angle DAB$ is a right angle. [3]
- (iii) Calculate the area of the quadrilateral $ABCD$. [2]

End of Paper

Answer key

1(i)	$\frac{20}{3}[3-2x]$ or $\frac{20}{3}[2x-3]$	7(ii)	$y = \frac{5}{4}x - \frac{17}{32}$
1(ii)	$x = 2\frac{2}{17}$ or 12 (NA)	7(iii)	Show $\frac{dy}{dx} > 0$ for $x > \frac{7}{16}$.
2	$a = 6, b = 10$	8(i)	$\frac{7}{25}$
3(a)	$x = 1.16$	8(ii)	$\frac{k}{\sqrt{1-k^2}}$
3(b)	$\frac{x}{y} = 3$	8(iii)	$\frac{4}{5}\sqrt{1-k^2} - \frac{3}{5}k$
4(i)	Constant term = $20a^3$	9(i)	$x = 1$
4(ii)	A term exists in $\frac{1}{x^3}$.	9(iii)	$(0, -3)$ and $(-6, 5)$
5(a)	$x = 0$ or $-\frac{8}{9}$ (NA)	9(iv)	AB is a diameter of the circle.
5(b)	$AD^2 = 51 - 6\sqrt{3}$	10(i)	-2 m/s^2
6(a)	$\frac{1}{2}\left(1 - \frac{3}{1-3x}\right)$ OR $\frac{2+3x}{2(3x-1)}$	10(ii)	$s = \frac{t^2}{2} - 7t + 12 \ln t+1 $
6(bi)	9	10(iii)	4.63 m
6(bii)	7	11(i)	$D(14, 11), M(14, 0), B(3, -7)$
6(biii)	$h = 8$	11(ii)	$\angle DAB$ is a right angle
		11(iii)	210 units ²

- 5 (a) The Population White Paper released by the Government of Singapore in 2013, projected that Singapore's population will hit 6.9 million by year 2030. The population of Singapore, P , increased from 5.399 million to 5.535 million from 2013 to 2015. Given that $P = Ae^{kt}$, where A and k are constants and t is the time in years from 2013.
- (i) Find the value of A and of k . [3]
- If the population continues to increase at the same rate,
- (ii) determine if the population trajectory for year 2030 in the Population White Paper is accurate. [2]

6 (i) Factorise completely the cubic polynomial $x^3 - x^2 + 3x - 3$. [2]


(ii) Express $\frac{2x^3 - 5x^2 + 10x - 3}{x^3 - x^2 + 3x - 3}$ in partial fractions. [5]

(iii) Differentiate $\ln(x^2 + 3)$ with respect to x . Hence express $\int_2^5 \frac{2x^3 - 5x^2 + 10x - 3}{x^3 - x^2 + 3x - 3} dx$ in the form $a + b \ln 2$, where a and b are integers. [5]

-

- (i) DE is parallel to BC , [1]
- (ii) $\triangle FDE$ is congruent to $\triangle EBF$, [3]
- (iii) $DEBF$ is a parallelogram. [2]

- 16/S4PR2/AM/2



The diagram shows the vertical cross-section $PQIRST$ of a structure, consisting of a triangle QRS of height 30 m and a rectangle $PQST$. The structure rests with PT on horizontal ground. To hold the structure up, a 15 m rope is secured at Q to a point, A , on the ground. It is given that QA is inclined at an angle, θ radians, to QP and $PT = 60 \sin \theta$ m.

- Show that the area, $A \text{ m}^2$, of the cross-section $PQIRST$ is given by

$$A = 900 \sin \theta + 450 \sin 2\theta.$$
 [4]
- Given that θ can vary, find the value of θ for which the maximum amount of paint is required to colour this cross-section. [5]
- Hence, find the maximum value of A . [1]

- 10 A trapezium of area, $A \text{ cm}^2$, has parallel sides of length $px^2 \text{ cm}$ and $q \text{ cm}$ and its perpendicular height is $x \text{ cm}$. Corresponding values of x and A are shown in the table below.


x	1	2	3	4
A	1.75	5	11.25	22

- Using suitable variables, draw, on a graph paper, a straight line graph and hence estimate the value of each of the constants p and q . [6]
- Using your values of p and q , calculate the value of x for which the trapezium is a rectangle. [2]
- Explain how another straight line drawn on your diagram can lead to an estimate of the value of x for which the trapezium is a rectangle. Draw this line and hence verify your value of x found in part (ii). [3]

- 11 Gravitational potential energy, measured in kilojoules (kJ), is the energy a body has due to its position. It can be calculated by the following equation:

$$\text{Gravitational potential energy} = \frac{mgh}{1000}$$

where m is the mass of the body in kg, g is the gravitational field strength in N/kg and h is the height of the object in m. The gravitational field strength, g , on Earth is approximately 10 N/kg.



The gravitational potential energy, E , in kJ, of a pirate ship ride can be modelled by the equation, $E = 100(1 - \cos kt) + a$, where k and a are constants and t is the time in seconds after starting the ride at loading level.

- Given that the mass of the ride is 1000 kg and at an initial loading level of 3 m, show that $a = 30$. [1]
 - Explain why this model suggests that the maximum gravitation potential energy possessed by the ride is 230 kJ. [1]
- The ride takes 6 seconds to travel from one peak to another.
- Show that the value of k is $\frac{\pi}{3}$ radians per second. [2]
 - Calculate the gravitation potential energy of the ride at $t = 8 \text{ s}$. [2]
 - If the ride continues for 60 seconds, find the exact duration for which the ride possesses more than 80 kJ of gravitational potential energy. [5]

End of Paper

This document is intended for internal circulation in Victoria School only. No part of this document may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying or otherwise, without the prior permission of the Victoria School Internal Exams Committee.

1	i	$a = 2$	8	ii	$2.24 \sin(x - 0.464) + 1$
	ii	$\left(-\frac{1}{2}, -1\frac{1}{2}\right)$		iii	a) Greatest value of $f(x) = 1 + \sqrt{5}$ Least value of $f(x) = 1 - \sqrt{5}$ b) Amplitude of $f(x) = \sqrt{5}$
2	a	$k \leq -9.32$ or $k \geq 3.32$	9	i	Show $QP = 15 \cos \theta$
	bi	$\frac{1}{4}$		ii	$\frac{\pi}{3}$
	ii	$1\frac{27}{64}$		iii	1170 m^2
3	i	Show base of similar pyramid $= \frac{2}{3}h$	10	i	Plot or $\frac{2A}{x} = px^2 + q$ $\frac{A}{x} = \frac{1}{2}px^2 + \frac{1}{2}q$ $p = 0.500$ $q = 3$
	ii	-1.35 cm s^{-1}		ii	$x = 2.45$
4	i	Let $f\left(\frac{1}{4}\right) = 0$ and $f(-2) = 0$		iii	Draw or $\frac{2A}{x} = x^2$ $\frac{A}{x} = 0.5x^2$ or $\frac{2A}{x} = 6$ or $\frac{A}{x} = 3$ $x = 2.45$
	ii	5	11	i	Sub. $m = 1000, g = 10, h = 3$
5	ai	$A = 5.399 \times 10^6$ $k = 0.0124$		ii	Sub. $\cos kt = -1$
	ii	Show $t = 17, P = 6.67 \times 10^6$ or $P = 6.9 \times 10^6, t = 19.72, \text{ Year} = 2032$ Not accurate		iii	Period $= \frac{2\pi}{k} = 6$
	b	4.77 units^2	iv	iv	180 kJ
6	i	$(x-1)(x^2+3)$		v	40 s
	ii	$2 + \frac{1}{x-1} - \frac{4x}{x^2+3}$	7	bi	Mid-Point Theorem
7	iii	$\frac{2x}{x^2+3}; 6 - 2 \ln 2$		ii	$\triangle FDE \equiv \triangle EBF$ (AAS)
8	i		8	i	